

Mediating effect of waist:height ratio on the association between BMI and frailty: the Korean Frailty and Aging Cohort Study

Minseo Kim^{1†}, Yunhwan Lee^{2†}, Eun-Young Kim¹ and Yongsoon Park^{1*}

¹Department of Food and Nutrition, Hanyang University, Seoul, Republic of Korea

²Department of Preventive Medicine and Public Health, Ajou University School of Medicine, Suwon, Republic of Korea

(Submitted 22 March 2019 – Final revision received 11 July 2019 – Accepted 6 August 2019 – First published online 27 August 2019)

Abstract

Body weight is a major risk determinant of frailty, but the effect of obesity on frailty is controversial. The present study aimed to confirm the hypothesis that the risk of frailty is positively associated with obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$), but the association is mediated by the waist:height ratio (WHtR) in older women and men. A total of 2862 community-dwelling older individuals aged 70–84 years were assessed for frailty using the Korean version of Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight index. Obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$) was associated with a higher risk of frailty compared with $\text{BMI} 18.5\text{--}<23 \text{ kg/m}^2$ in all the older individuals (OR 1.88; 95 % CI 1.11, 3.17; $P = 0.018$) and in older women (OR 1.86; 95 % CI 1.01, 3.42; $P = 0.047$) before adjusting for WHtR but was not associated with BMI after adjusting for WHtR. Additionally, obesity was not significantly associated with the risk of frailty before and after adjusting for WHtR in older men. Mediation analysis revealed that the association between BMI and frailty score was mediated by WHtR. Moreover, the mediating effect of WHtR on frailty score was positive in both women and men, but the frailty score was associated with BMI positively in women and negatively in men. The present study suggests that the risk of frailty is higher in obese women, which is mediated by WHtR, but not in obese men.

Key words: BMI: Older adults: Frailty: Mediation analysis: Waist:height ratio

Frailty is a state of high vulnerability for adverse health outcomes, including disability, dependency, falls, need for long-term care and mortality⁽¹⁾. Korea is one of the most rapidly ageing countries worldwide⁽²⁾, and the prevalence of frailty defined by the Korean version of the Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight (K-FRAIL) index has been reported to be 12.4 % in Koreans aged 70 years or older⁽³⁾.

One of the important factors for frailty is weight loss since the decline in body weight leads to frequently observed underweight in the older adults aged 70 years or older^(4,5). Epidemiological studies consistently reported that the prevalence of frailty was higher in the older adults who are underweight ($\text{BMI} < 20 \text{ kg/m}^2$)^(6–10) and lower in those who are overweight ($\text{BMI} 25\text{--}29.9 \text{ kg/m}^2$)^(8–10). However, the association between obesity and frailty is controversial. The previous studies showed that obesity defined by $\text{BMI} \geq 30 \text{ kg/m}^2$ was positively associated with the prevalence and incidence of frailty in the older adults^(8,10–16), but a few studies showed no significant association between obesity and the risk of frailty^(7,9,16).

Garcia-Esquinas *et al.*⁽¹⁶⁾ reported that obesity was positively associated with the risk of frailty in the older adults, but the association was not significant after adjusting for waist circumference (WC), suggesting that the association between BMI and frailty could be mediated by abdominal obesity. A few previous studies reported that abdominal obesity was associated with higher prevalence and incidence of frailty in the older adults^(8,9,16,17). Increased WC has been suggested to be associated with a marker of oxidative stress and inflammation, independent of BMI⁽¹⁸⁾, which may be the major mechanism leading to frailty⁽¹⁹⁾.

Abdominal obesity is commonly defined by WC, and waist:height ratio (WHtR) has been suggested to be a better predictor of whole-body fat percentage and visceral adipose tissue mass as well as increased health risk compared with WC⁽²⁰⁾. Partezani-Rodrigues *et al.*⁽²¹⁾ reported that in the older adults the risk of frailty was significantly associated with WHtR but not with WC. However, no study investigated the mediating effect of WHtR on the association between BMI and the risk of frailty. Therefore, the present study aimed to confirm the hypothesis

Abbreviations: K-FRAIL, Korean version of the Fatigue, Resistance, Ambulation, Illnesses, and Loss of weight; WC, waist circumference; WHtR, waist:height ratio.

* **Corresponding author:** Professor Yongsoon Park, fax +82-2220-1856, email yongsoon@hanyang.ac.kr

† These authors contributed equally to this work.



that the risk of frailty was positively associated with obesity, but the association was mediated by WHtR in the older women and men.

Methods

Participants

The present study used the data from the Korean Frailty and Aging Cohort Study (KFACS), a multicentre, longitudinal study of community-dwelling older adults⁽²²⁾. Between June 2016 and November 2017, the participants of the KFACS were recruited from ten participating centres across the urban and rural areas of Korea on the basis of age- and sex-specific strata. The following participants were included in the study: those aged 70–84 years, those living independently at home, those with no plans to move out in the next 2 years and those having no serious problems with communication. Among the 3014 participants who completed the initial assessment, 2997 participants with non-missing data on frailty assessments were included in the present study. The study was conducted according to the guidelines in the Declaration of Helsinki⁽²³⁾ and was approved by the institutional review board of Kyung Hee University (KHUH-2015-12-103-044) and the institutional review board of Hanyang University (HYI-18-167-1). All participants signed a written informed consent form.

Data collection

Data were obtained from the KFACS⁽²²⁾ and included age, sex, height, weight, WC, educational level (less than middle school or more than high school), job (currently working, previously working or non-working), smoking (current, former or never), alcohol drinking (once or more times in a month) and living alone. Polypharmacy was defined as the regular use of five or more prescription medications. Cognitive impairment was defined as a Korean Mini-Mental State Examination score of less than 24⁽²⁴⁾. Co-morbidity was defined as the number of self-reported diseases diagnosed by a physician (hypertension, dyslipidaemia, CVD, peripheral arterial disease, cerebrovascular disease, osteoarthritis, osteoporosis, respiratory disease, lung disease, thyroid disease, dementia, kidney disease and cancer). Disabilities were defined as dependence in at least one item in the Korean activities of daily living⁽²⁵⁾ and Korean instrumental activities of daily living⁽²⁶⁾.

Weight was measured to the nearest 0.1 kg using a portable digital scale, and height was measured to the nearest 0.1 cm using a measuring tape. The BMI was calculated as weight divided by square of the height. According to the criteria of the World Health Organization Asia-Pacific Region⁽²⁷⁾, BMI was categorised as <18.5 kg/m², 18.5–<23 kg/m², 23–<25 kg/m², 25–<30 kg/m² and ≥30 kg/m². The WC was measured at the middle point between the rib margin and iliac crests in a horizontal plane using an inelastic measuring tape. The WHtR (cm/cm) was calculated as WC divided by height with the median value of 0.557 as the cut-off point in the study participants. Dual-energy X-ray absorptiometry was used to obtain appendicular skeletal muscle mass (ASM) of the four limbs, and ASM index was

calculated as the sum of muscle mass in the four limbs (kg) divided by the squared height (m²).

Frailty indices

Frailty was defined using the K-FRAIL index⁽²⁸⁾, a modified version of the published FRAIL scale⁽²⁹⁾. The FRAIL scale exhibited the strongest predictive validity for disability and mortality compared with the Cardiovascular Health Study (CHS) index⁽³⁰⁾. Since the K-FRAIL index is easier to measure compared with CHS because of interview-based test⁽³¹⁾, the K-FRAIL index was measured in all participants in the study. The K-FRAIL index assigns 1 point to each of the following five components. Fatigue was assessed by asking the participants for how much time during the preceding 4 weeks they felt tired, with responses of 'all of the time' or 'most of the time' scored as 1 point. Resistance was defined as difficulty in walking up to ten stair steps alone without resting and without aid, and ambulation was assessed by asking whether they had any difficulty in walking 300 m alone and without aid. Illness was defined as having five or more conditions among the following eleven conditions: hypertension, diabetes mellitus, chronic obstructive pulmonary disease, angina, myocardial infarction, heart failure, asthma, arthritis, stroke, renal disease and cancer. Loss of weight was recorded as loss of at least 5 % of the body weight within the preceding year. Participants with a total score of 3 or higher are classified as frail.

Statistical analysis

Independent two-sample *t* tests for continuous variables and χ^2 tests for categorical variables were used to compare the differences in the characteristics and risk factors for frailty between older women and men. Continuous variables were presented as means and standard deviations, and categorical variables were expressed as the number of participants (percentage distribution). Pearson's correlation analysis was performed to determine the bivariate association between the main variables. To assess whether the association between BMI and risk of frailty weakened or became non-significant after adjusting for WHtR, binary logistic regression analysis was performed after adjusting for confounding factors. ANCOVA with least significant difference *post hoc* test was used in comparing the marginal means of lean mass by adjusting the confounding variables.

Moderation, mediation and moderated mediation analyses were performed using the Hayes PROCESS macro (models 1, 4 and 14)⁽³²⁾. For these analyses, independent, dependent and mediator variables or categorical covariates were designated as numerical variables or dummy variables. Additionally, moderator variables were recoded using indicator coding, with men as the reference category⁽³³⁾. This macro used a bootstrapping strategy to test the validity of the indirect effects and calculate the 95 % bias-corrected CI from 5000 bootstrap samples⁽³⁴⁾. These analyses were considered significant if the CI excluded zero. In the multivariable models, the covariates showing a significance level below 0.2 were included in the final model⁽³⁵⁾. All statistical tests were two sided according to a 0.05 significance level and performed using SPSS (version 25.0; SPSS Inc.).

Table 1. Characteristics of women and men*
(Mean values and standard deviations; percentages)

	Total (n 2862)		Women (n 1482)		Men (n 1380)		P†
	Mean	SD	Mean	SD	Mean	SD	
Age (years)	75.96	3.89	75.66	3.85	76.29	3.91	<0.001
Height (cm)	157.88	8.54	151.50	5.27	164.72	5.55	<0.001
Weight (kg)	61.03	9.33	57.26	7.93	65.08	9.00	<0.001
BMI (kg/m ²)	24.46	3.01	24.92	3.03	23.96	2.92	<0.001
BMI (%)							
<18.5 kg/m ²	1.9		0.9		3.0		<0.001
18.5–<23 kg/m ²	29.3		25.2		33.7		
23–<25 kg/m ²	28.5		28.5		28.6		
25–<30 kg/m ²	36.1		39.9		32.1		
≥30 kg/m ²	4.1		5.5		2.6		
WC (cm)	87.83	8.49	87.05	8.37	88.66	8.56	<0.001
WHtR (cm/cm)	0.56	0.06	0.58	0.06	0.54	0.05	<0.001
Education (%)							<0.001
<High school	62.3		76.0		47.7		
≥High school	37.7		24.0		52.3		
Currently working (%)	26.2		22.1		30.5		<0.001
Smoking (%)							<0.001
Current	5.7		0.9		10.9		
Former	33.7		2.0		67.7		
Never	60.6		97.1		21.4		
Alcohol drinking (%)	37.5		19.7		56.5		<0.001
Living alone (%)	22.4		35.2		8.7		<0.001
Polypharmacy (%)	33.0		30.3		35.9		0.001
Cognitive impairment (%)	20.9		26.5		15.0		<0.001
Co-morbidity (%)							<0.001
0	16.9		10.9		23.3		
1	26.2		22.5		30.1		
≥2	56.9		66.5		46.6		
ADL disability (%)	11.0		13.7		8.2		<0.001
IADL disability (%)	37.8		18.4		58.5		<0.001
Frailty (%)	12.3		17.1		7.2		<0.001

WC, waist circumference; WHtR, waist:height ratio; ADL, activities of daily living; IADL, instrumental activities of daily living.

* Number of participants was discordant; education (n 2861), at work (n 2860), smoking (n 2861), alcohol drinking (n 2852), polypharmacy (n 2848) and IADL (n 2857).

† P values were analysed using the independent t test for continuous variables and χ^2 test for categorical variables.

Results

Characteristics of older women and men

The prevalence of frailty was 12.3 % and was higher in women than in men (Table 1). Women were younger; and had lower educational level, proportion of currently working, smoking, alcohol drinking, polypharmacy and instrumental activities of daily living disability. However, women had higher BMI, WHtR, proportion of living alone, cognitive impairment, co-morbidity and activities of daily living disability compared with men.

Association between frailty and body composition

Multivariable-adjusted regression analysis revealed that the risk of frailty was higher in all older individuals and in older men with BMI < 18.5 kg/m² before and after adjusting for WHtR (Table 2). Furthermore, the risk of frailty was higher in all older individuals and in older women with BMI ≥ 30 kg/m² before adjusting for WHtR, but the risk disappeared after adjusting for WHtR. On the other hand, the risk of frailty was lower in men with BMI 25–<30 kg/m² before adjusting for WHtR but lower in all older individuals and older women with BMI 25–<30 kg/m² after adjusting for WHtR.

When participants were divided into low and high WHtR groups according to the median value of 0.557 as cut-off point, the risk of frailty was higher in all older individuals and in older men with BMI < 18.5 kg/m² and low WHtR compared with those with BMI 18.5–<23 kg/m² and low WHtR (Table 3). Moreover, the risk of frailty was higher in all older individuals and women with BMI ≥ 30 kg/m² and high WHtR than those with BMI 18.5–<23 kg/m² and low WHtR. On the other hand, the risk of frailty was lower in men with BMI 25–<30 kg/m² and high WHtR compared with those with BMI 18.5–<23 kg/m² and low WHtR.

Mediating effects of waist:height ratio on the association between BMI and frailty

Bivariate correlations revealed that BMI was positively correlated with WHtR (r 0.82, P < 0.001) and frailty (r 0.04, P = 0.024), and WHtR was positively correlated with frailty (r 0.18, P < 0.001) (online Supplementary Table S1). In the mediation analysis, the negative direct effect and the positive indirect effect of BMI on the score of frailty were significant in all older individuals, in older women and in older men (Fig. 1). On the other hand, the total effect of BMI on the score of frailty was positive in women but negative in men. Additionally,



Table 2. Association between BMI and risk of frailty before and after adjusting for waist:height ratio (WHtR)* (Odds ratios and 95 % confidence intervals)

BMI (kg/m ²)	Model 1						Model 2					
	Total		Women		Men		Total		Women		Men†	
	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI
<18.5	3.19	1.55, 6.57	1.81	0.48, 6.79	4.19	1.73, 10.17	4.02	1.88, 8.57	2.33	0.60, 9.05	3.74	1.36, 10.26
18.5–<23	1.0		1.0		1.0		1.0		1.0		1.0	
23–<25	1.02	0.73, 1.41	0.92	0.61, 1.40	1.10	0.64, 1.88	0.86	0.60, 1.24	0.72	0.45, 1.14	1.19	0.63, 2.22
25–<30	0.76	0.55, 1.06	0.92	0.62, 1.36	0.40	0.21, 0.75	0.56	0.36, 0.87	0.58	0.34, 0.99	0.45	0.19, 1.07
≥30	1.88	1.11, 3.17	1.86	1.01, 3.42	1.89	0.64, 5.57	1.08	0.51, 2.28	0.82	0.34, 1.96	2.47	0.51, 11.97

ADL, activities of daily living; IADL, instrumental activities of daily living.

* For logistic regression model 1, it was adjusted for age, sex, education, alcohol drinking, polypharmacy, cognitive impairment, co-morbidity, ADL and IADL in total older individuals (*n* 2830); age, education, polypharmacy, cognitive impairment, co-morbidity, ADL and IADL in women (*n* 1465); age, education, alcohol drinking, living alone, polypharmacy, co-morbidity and IADL in men (*n* 1365). For logistic regression model 2, model 1 adjusted additionally for WHtR in total older adults, women and men.

† As WHtR was a NS covariate variable (*P* = 0.643), the Hosmer–Lemeshow test was significant (χ^2 = 16.689, *df* = 8, *P* = 0.034) for men after adjusting for WHtR in model 2.

Table 3. Subgroup analysis of waist:height ratio (WHtR) on the association between BMI and risk of frailty*† (Odds ratios and 95 % confidence intervals)

	BMI (kg/m ²)									
	<18.5		18.5–<23		23–<25		25–<30		≥30	
	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI
Total (<i>n</i> 2830)										
Low WHtR	3.32	1.61, 6.86	1.0		1.08	0.70, 1.67	0.29	0.10, 0.82	N/A	
High WHtR	N/A		1.41	0.74, 2.67	1.07	0.71, 1.61	0.88	0.62, 1.24	2.00	1.17, 3.42
Women (<i>n</i> 1465)										
Low WHtR	2.08	0.55, 7.87	1.0		0.76	0.40, 1.46	0.70	0.23, 2.13	N/A	
High WHtR	N/A		1.74	0.86, 3.50	1.21	0.74, 1.98	1.08	0.70, 1.67	2.13	1.13, 4.01
Men (<i>n</i> 1365)										
Low WHtR	4.16	1.70, 10.18	1.0		1.26	0.70, 2.27	N/A		N/A	
High WHtR	N/A		N/A		0.72	0.32, 1.64	0.50	0.26, 0.95	1.84	0.62, 5.43

N/A, not applicable; ADL, activities of daily living; IADL, instrumental activities of daily living.

* N/A was used if there were no frail or older adult participants.

† Confounding factors were age, sex, education, alcohol drinking, polypharmacy, cognitive impairment, co-morbidity, ADL and IADL in total older adults; age, education, polypharmacy, cognitive impairment, co-morbidity, ADL and IADL in women; age, education, alcohol drinking, living alone, polypharmacy, co-morbidity, ADL and IADL in men.

the mediating effect of WHtR on the score of frailty was positive in both women and men, but the score of frailty was associated with BMI positively in women but negatively in men, suggesting that the association was affected by sex (online Supplementary Fig. S1).

Discussion

The present study showed that the risk of frailty was higher in the older individuals with BMI ≥ 30 kg/m² than those with BMI 18–<23 kg/m² before adjusting for WHtR but not after adjusting for WHtR in total older individuals and women but not in men. The association between BMI and the risk of frailty was mediated by WHtR in the older individuals, particularly in women.

Consistent with the present study, the majority of previous studies reported that obesity (BMI ≥ 30 kg/m²) was associated with a higher prevalence or incidence of frailty in the older adults without adjusting for abdominal obesity^(8,10–16). However, a few studies showed that the risk of frailty was not significantly higher in obese older adults^(7,9,16). This inconsistency could be partly because of the proportion of frailty among the female study participants. Older women tended to be more prone to obesity⁽³⁶⁾

and had a higher prevalence of frailty compared with older men⁽³⁷⁾. In previous studies reporting the absence of an association between obesity and frailty, the proportion of frailty in women (8–9 %) was lower than that in the present study (17 %)^(7,9). Oestrogen levels declined after menopause, leading to a decrease in bone density, muscle mass and muscle strength but an increase in visceral adiposity⁽³⁸⁾. Consistent with the findings of the present study, the prevalence of obesity was higher in women than in men⁽³⁹⁾. Additionally, previous studies did not include abdominal obesity as a confounding factor to confirm the association between frailty and obesity^(7–15). In the present study, obesity was also associated with a higher risk of frailty before adjusting for WHtR, but it was not associated with the risk of frailty after adjusting for WHtR. Previous epidemiological studies consistently reported that abdominal obesity defined by WC^(8,9,16,17,40) and WHtR⁽²¹⁾ was associated with a higher prevalence and incidence of frailty in the older adults. Older adults with high WC had high levels of oxidative stress markers, independent of BMI, suggesting a possible involvement of oxidative stress in the genesis of frailty^(18,19). Garcia-Esquinas *et al.*⁽¹⁶⁾ reported that the association between obesity and the risk of frailty was not significant in the older adults after adjusting




Fig. 1. Mediating effects of waist:height ratio (WHtR) on the association between BMI and frailty score in total older adults (A), women (B) and men (C). Unstandardised coefficients and 95 % confidence intervals are presented: 'a' is the linear regression coefficient of the BMI–WHtR association and 'b' is that of the WHtR–frailty association. Adjusted confounding factors were age, sex, education, alcohol drinking, polypharmacy, cognitive impairment, activities of daily living (ADL) and instrumental activities of daily living (IADL) in total older adults; age, education, polypharmacy, cognitive impairment, ADL and IADL in women; age, education, alcohol drinking, living alone, polypharmacy, ADL and IADL in men. * $P < 0.05$, ** $P < 0.001$.

for WC, suggesting that the association could be mediated by abdominal obesity. Moreover, Ferrioli *et al.*⁽⁹⁾ suggested that the WC may be one of the main factors that associate obesity with frailty in the older adults. However, Rodrigues *et al.*⁽²¹⁾ reported that the risk of frailty was positively associated with WHtR ≥ 0.50

but not with WC, suggesting that WHtR could be a better marker of abdominal obesity. Hirani *et al.*⁽⁴¹⁾ also reported that the risk of frailty was significantly higher in the older adults with sarcopenic obesity than in those with non-sarcopenic normal weight, but it was not significantly different from those with



non-sarcopenic obesity. It is well known that sarcopenia, defined as an abnormal loss of muscle mass and strength, is one of the major causes for an increase in the risk of frailty⁽⁴²⁾. Falsarella *et al.*⁽⁴³⁾ reported that frail older individuals had a lower muscle mass and higher percentage of body fat compared with non-frail older individuals. In the present study, the obese older individuals had a higher lean mass compared with the older individuals with normal weight (online Supplementary Table S2).

In the present study, obesity was not associated with risk of frailty in men before and after adjusting for WHtR. The mediating effect of WHtR on the association between BMI and risk of frailty was shown only in women but not in men, suggesting that the mediating effect was moderated by sex (online Supplementary Fig. S1). In the present study, it could be impossible to detect the association between BMI and the risk of frailty in men, because there were only five frail men with obesity. Furthermore, according to the previous studies, men had a higher muscle mass compared with women⁽⁴⁴⁾, and high muscle mass has been shown to be associated with lower risk of frailty in the older adults⁽⁴⁵⁾. The present study also showed that men had more muscle mass compared with women (online Supplementary Table S2). Bigaard *et al.*⁽⁴⁶⁾ reported that WC reflected visceral fat, whereas BMI reflected fat-free mass or fat deposit elsewhere, suggesting that BMI in men could not be associated with the risk of frailty because of the body composition effect. In the present study, the effect of BMI on frailty score was positive in women, but negative in men, suggesting that high muscle mass in men could attenuate the adverse effect of obesity through WHtR on the risk of frailty (online Supplementary Fig. S1).

The previous studies consistently suggested that underweight was associated with higher prevalence and incidence of frailty^(6–10, 12–15). Underweight could have resulted from malnutrition, cachexia and sarcopenia, which were also associated with increased risk of frailty and represent low reserve capacity and weight loss, a key component of frailty⁽⁴⁷⁾. In the present study, underweight was associated with higher risk of frailty in the older men before and after adjusting for WHtR, suggesting that WHtR was not associated with the risk of frailty in men who are underweight. However, in the present study, the risk of frailty was not higher in underweight women because of a small proportion of underweight women with frailty (n 4).

Previous studies reported that overweight defined by BMI 25–29.9 kg/m² was associated with lower prevalence and incidence of frailty in the older adults^(8–10, 17). Overweight was also associated with lower risk of mortality in the older adults, suggesting that overweight could represent a higher reserve capacity than those who are underweight or even normal weight⁽⁷⁾. Oreopoulos *et al.*⁽⁴⁸⁾ reported that a higher BMI in advanced age can actually be considered a protective factor against oxidative stress, inflammation, malnutrition, fractures and cognitive decline. Additionally, the negative association between BMI and mortality suggested a paradoxical association

of overweight or obesity with reduced CVD-related mortality in the older adults⁽⁴⁹⁾. In the present study, BMI 25–29.9 kg/m² was associated with lower risk of frailty in all older individuals and in women after adjusting for WHtR.

The present study has a few limitations. First, the present study was composed of only a small proportion of frail older individuals. Second, the cross-sectional study design was not able to identify the causal association between BMI and WHtR with the risk of frailty. Third, although adjustments for confounders were made, unmeasured factors might affect the results of the present study.

In conclusion, the present study showed that the risk of frailty was higher in obese individuals, which was mediated by WHtR in older individuals, particularly older women, suggesting that abdominal obesity could increase the risk of frailty in women but not in men. A large population-based prospective observational study is needed to confirm whether the reduction in WHtR in obese older adults reduces the risk of frailty.

Acknowledgements

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean Government (MSIT) (NRF-2018R1A2B6002486) and a grant of the Korea Health Technology R&D Project through the Korean Health Industry Development Institute (KHIDI) funded by the Ministry of Health & Welfare, the Republic of Korea (grant number: HI15C3153).

The authors' contributions are as follows: M. K. performed the statistical analyses and prepared the first draft, Y. L. interpreted the results and edited the manuscript, E. K. assisted with statistical analyses and interpretation of results and edited the manuscript and Y. P. designed the study and finalised the manuscript. All authors read and approved the final manuscript.

The authors declare that there are no conflicts of interest.

Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114519002058>

References

1. Fried LP, Ferrucci L, Darer J, *et al.* (2004) Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci* **59**, M255–M263.
2. He W, Goodkind D & Kowal P (2016) An aging world: 2015. International population reports. Washington DC: U.S. Census Bureau.
3. Kim KJ, Shin J, Choi J, *et al.* (2018) Discrepancies in the prevalence of known frailty scales: Korean Frailty and Aging Cohort Study. *Ann Geriatr Med Res* **22**, 137–144.
4. Morley JE (2007) Weight loss in older persons: new therapeutic approaches. *Curr Pharm Des* **13**, 3637–3647.

5. Reinders I, Visser M & Schaap L (2017) Body weight and body composition in old age and their relationship with frailty. *Curr Opin Clin Nutr Metab Care* **20**, 11–15.
6. Sanchez-Garcia S, Sanchez-Arenas R, Garcia-Pena C, *et al.* (2014) Frailty among community-dwelling elderly Mexican people: prevalence and association with sociodemographic characteristics, health state and the use of health services. *Geriatr Gerontol Int* **14**, 395–402.
7. Lee Y, Kim J, Han ES, *et al.* (2014) Frailty and body mass index as predictors of 3-year mortality in older adults living in the community. *Gerontology* **60**, 475–482.
8. Hubbard RE, Lang IA, Llewellyn DJ, *et al.* (2010) Frailty, body mass index, and abdominal obesity in older people. *J Gerontol A Biol Sci Med Sci* **65**, 377–381.
9. Ferrioli E, Pessanha F, Moreira VG, *et al.* (2017) Body composition and frailty profiles in Brazilian older people: Frailty in Brazilian Older People Study-FIBRA-BR. *Arch Gerontol Geriatr* **71**, 99–104.
10. Rietman ML, Van Oostrom S, Picavet H, *et al.* (2018) The association between BMI and different frailty domains: a U-shaped curve? *J Nutr Health Aging* **22**, 8–15.
11. Sheehan KJ, O'Connell MD, Cunningham C, *et al.* (2013) The relationship between increased body mass index and frailty on falls in community dwelling older adults. *BMC Geriatr* **13**, 132.
12. Lopez SR, Montero P & Carmenate M (2014) Educational inequalities and frailty in Spain: what is the role of obesity? *J Frailty Aging* **3**, 120–125.
13. Woods NF, LaCroix AZ, Gray SL, *et al.* (2005) Frailty: emergence and consequences in women aged 65 and older in the Women's Health Initiative Observational Study. *J Am Geriatr Soc* **53**, 1321–1330.
14. Blaum CS, Xue QL, Tian J, *et al.* (2009) Is hyperglycemia associated with frailty status in older women? *J Am Geriatr Soc* **57**, 840–847.
15. Blaum CS, Xue QL, Michelson E, *et al.* (2005) The association between obesity and the frailty syndrome in older women: the Women's Health and Aging Studies. *J Am Geriatr Soc* **53**, 927–934.
16. Garcia-Esquinas E, Jose Garcia-Garcia F, Leon-Munoz LM, *et al.* (2015) Obesity, fat distribution, and risk of frailty in two population-based cohorts of older adults in Spain. *Obesity (Silver Spring)* **23**, 847–855.
17. Liao Q, Zheng Z, Xiu S, *et al.* (2018) Waist circumference is a better predictor of risk for frailty than BMI in the community-dwelling elderly in Beijing. *Aging Clin Exp Res* **30**, 1319–1325.
18. Weinbrenner T, Schröder H, Escuriol V, *et al.* (2006) Circulating oxidized LDL is associated with increased waist circumference independent of body mass index in men and women. *Am J Clin Nutr* **83**, 30–35.
19. Serviddio G, Romano A, Greco A, *et al.* (2009) Frailty syndrome is associated with altered circulating redox balance and increased markers of oxidative stress. *Int J Immunopathol Pharmacol* **22**, 819–827.
20. Swainson MG, Batterham AM, Tsakirides C, *et al.* (2017) Prediction of whole-body fat percentage and visceral adipose tissue mass from five anthropometric variables. *PLOS ONE* **12**, e0177175.
21. Rodrigues RP, Fhon JS, Huayta VR, *et al.* (2017) Frailty syndrome and anthropometric measurements in the elderly living at home. *J Aging Res Clin Pract* **6**, 133–138.
22. Won CW, Lee Y, Choi J, *et al.* (2016) Starting construction of frailty cohort for elderly and intervention study. *Ann Geriatr Med Res* **20**, 114–117.
23. Dale O & Salo M (1996) The Helsinki Declaration, research guidelines and regulations: present and future editorial aspects. *Acta Anaesthesiol Scand* **40**, 771–772.
24. Kang Y, Na DL & Hahn S (1997) A validity study on the Korean Mini-Mental State Examination (K-MMSE) in dementia patients. *J Korean Neurol Assoc* **15**, 300–308.
25. Won CW, Rho YG, Kim SY, *et al.* (2002) The validity and reliability of Korean Activities of Daily Living (K-ADL) scale. *J Korean Geriatr Soc* **6**, 98–106.
26. Won CW, Rho YG, Duk S, *et al.* (2002) The validity and reliability of Korean Instrumental Activities of Daily Living (K-IADL) scale. *J Korean Geriatr Soc* **6**, 273–280.
27. World Health Organization (2000) *The Asia-Pacific Perspective: Redefining Obesity and Its Treatment*. Sydney: Health Communications Australia.
28. Jung HW, Yoo HJ, Park SY, *et al.* (2016) The Korean version of the FRAIL scale: clinical feasibility and validity of assessing the frailty status of Korean elderly. *Korean J Intern Med* **31**, 594–600.
29. Morley JE, Malmstrom T & Miller D (2012) A simple frailty questionnaire (FRAIL) predicts outcomes in middle aged African Americans. *J Nutr Health Aging* **16**, 601–608.
30. Malmstrom TK, Miller DK & Morley JE (2014) A comparison of four frailty models. *J Am Geriatr Soc* **62**, 721–726.
31. Woo J, Leung J & Morley JE (2012) Comparison of frailty indicators based on clinical phenotype and the multiple deficit approach in predicting mortality and physical limitation. *J Am Geriatr Soc* **60**, 1478–1486.
32. Hayes AF (2013) *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-based Approach*. New York: The Guilford Press.
33. Hayes AF & Preacher KJ (2014) Statistical mediation analysis with a multicategorical independent variable. *Br J Math Stat Psychol* **67**, 451–470.
34. Hayes AF (2009) Beyond Baron and Kenny: statistical mediation analysis in the new millennium. *Commun Monogr* **76**, 408–420.
35. Greenland S & Pearce N (2015) Statistical foundations for model-based adjustments. *Annu Rev Public Health* **36**, 89–108.
36. Ogden CL, Yanovski SZ, Carroll MD, *et al.* (2007) The epidemiology of obesity. *Gastroenterology* **132**, 2087–2102.
37. Collard R, Boter H, Schoevers R, *et al.* (2012) Prevalence of frailty in community-dwelling older persons: a systematic review. *J Am Geriatr Soc* **60**, 1487–1492.
38. Messier V, Rabasa-Lhoret R, Barbat-Artigas S, *et al.* (2011) Menopause and sarcopenia: a potential role for sex hormones. *Maturitas* **68**, 331–336.
39. Hong KS, Bang OY, Kang DW, *et al.* (2013) Stroke statistics in Korea: part I. Epidemiology and risk factors: a report from the Korean stroke society and clinical research center for stroke. *J Stroke* **15**, 2–20.
40. Liu LK, Lee WJ, Chen LY, *et al.* (2015) Association between frailty, osteoporosis, falls and hip fractures among community-dwelling people aged 50 years and older in Taiwan: results from I-Lan Longitudinal Aging Study. *PLOS ONE* **10**, e0136968.
41. Hirani V, Naganathan V, Blyth F, *et al.* (2017) Longitudinal associations between body composition, sarcopenic obesity and outcomes of frailty, disability, institutionalisation and mortality in community-dwelling older men: The Concord Health and Ageing in Men Project. *Age Ageing* **46**, 413–420.
42. Yu Z, Ruan Q, & D'Onofrio G (2017) From Sarcopenia to Frailty: the pathophysiological basis and potential target molecules of intervention. In *Frailty and Sarcopenia-Onset, Development and Clinical Challenges*, pp. 56–57 [Y Dionysiotis, editor]. Croatia: InTech.
43. Falsarella GR, Gasparotto LPR, Barcelos CC, *et al.* (2015) Body composition as a frailty marker for the elderly community. *Clin Interv Aging* **10**, 1661–1666.
44. Bredella MA (2017) Sex differences in body composition. In *Sex and Gender Factors Affecting Metabolic Homeostasis, Diabetes and Obesity*, p. 12 [FM Jarvis, editor]. Basel, Switzerland: Springer.



45. Meng L, Shi H, Shi J, *et al.* (2019) Differences in clinical characteristics, muscle mass, and physical performance among different frailty levels in Chinese older men. *Chin Med J* **132**, 352–355.
46. Bigaard J, Tjønneland A, Thomsen BL, *et al.* (2003) Waist circumference, BMI, smoking, and mortality in middle-aged men and women. *Obes Res* **11**, 895–903.
47. Rolland Y, Van Kan GA, Gillette Guyonnet S, *et al.* (2011) Cachexia versus sarcopenia. *Curr Opin Clin Nutr Metab Care* **14**, 15–21.
48. Oreopoulos A, Kalantar-Zadeh K, Sharma AM, *et al.* (2009) The obesity paradox in the elderly: potential mechanisms and clinical implications. *Clin Geriatr Med* **25**, 643–659.
49. Kim TN (2018) Elderly obesity: is it harmful or beneficial? *J Obes* **27**, 84–92.